arXiv:0909.0109v2 [cs.AI] 22 Jan 2010

A Layered Graph Representation for Complex Regions

Sanjiang Li
Centre for Quantum Computation and Intelligent Systems,
Faculty of Engineering and Information Technology,
University of Technology, Sydney, Broadway NSW 2007, Australia
sanjiang.li@uts.edu.au

Abstract

Topological information are the most important kind of qualitative spatial information. Current formalisms for the topological aspect of space focus on relations between regions or properties of regions. This work provides a qualitative model for representing the topological internal structure of complex regions, which could be of multiple pieces and/or have holes and islands to any finite level. We propose a layered graph model for representing the internal structure of complex plane regions, where each node represents the closure of a connected component of the interior or the exterior of a complex region. The model provides a complete representation in the sense that the (global) nine-intersections between the interiors, the boundaries, and the exteriors of two complex regions can be determined by the (local) RCC8 topological relations between the associated simple regions. Moreover, this graph model has an inherent hierarchy which could be exploited for map generalization.

keywords: qualitative spatial reasoning, topological relation, internal structure, layered graph, hole, connected component, GIS

Introduction

Qualitative spatial reasoning (QSR) is an established research subfield in Artificial Intelligence and Geographical Information Science (GISc). One basic requirement of QSR is to provide qualitative, non-numerical information at various details for spatial representation and reasoning. QSR is an interdisciplinary research of computer science, cognition, and geography, with important applications in areas including Geographical Information Systems (GISs), Content-Based Image Retrieval, Computer Graphics, Robotics (Cohn and Renz 2007).

Among the many aspects of space, topology is the most important one. A major part of QSR research focuses on the study of topological relation models and topological properties. The Region Connection Calculus (RCC) (Randell, Cui, and Cohn 1992) is perhaps the most well-known logic-based approach to topological information. Among the many topological relations defined in the RCC, the eight RCC8 basic relations are identified as of great importance. Topological properties, such as a region '*has a hole*' or '*has up to k-components*' can also be expressed in the RCC and in similar logical-based formalisms (Cohn and Hazarika 2001; Varzi 2007; Kontchakov et al. 2008).

This paper will not propose new topological relation model, instead, we will focus on the internal topological structure of complex regions.

As far as topology is concerned, closed disks are the most simple regions. Regions that are topologically equivalent to a closed disk are called *simple regions*. The inadequacy of simple regions for representing spatial phenomena, *e.g.* countries, is clear. Various models have been proposed to represent complex spatial regions. These include simple region with holes (Egenhofer, Clementini, and Di Felice 1994), composite region (Clementini, Di Felice, and Califano 1995), the tree model of (Worboys and Bofakos 1993), and the component view of (Schneider and Behr 2006).

Simple regions with holes (srhs) and composite regions are two special kinds of bounded regions. A srh is a bounded connected region with several holes, and a composite region is a bounded but possibly disconnected region that has no holes. Schneider and Behr (Schneider and Behr 2006) considered bounded regions in general. They decomposed each bounded region A into finite *face*s, *i.e.* strong connected components of A.

The relative position between different faces is lost in this approach. For example, it does not distinguish the situation where one face is contained in a hole of another from the situation where one face is outside another. For example, the two regions in Fig. 1 both have two *faces* a_1, a_2, while a_1 is a simple region with a single hole, and a_2 is a simple region. This suggests inclusion information between faces is neglected. Moreover, the cavity surrounded by several faces are not expressed in this approach. For example, the two regions in Fig. 2 have the same faces, but the right region has a hole surrounded by the two faces, while the left region has no holes.

(Worboys and Bofakos 1993) proposed a tree model for representing complex regions. This model, called W-B tree model, reflects the hierarchical inclusion among holes and islands of a complex region to any finite level. Unlike the approach of (Schneider and Behr 2006), the tree model does not decompose complex regions into disjoint faces. Instead, it represents complex regions as "constrained and structured aggregations of atoms." As a result, it is irrelevant how the

---

Copyright © 2010, Association for the Advancement of Artificial Intelligence (www.aaai.org). All rights reserved.



Figure 1: Two complex regions with same faces



Figure 2: Two complex regions with same faces

holes separate their carrier in this model. For example, the two regions in Fig. 3 are represented by the same W-B tree.

In this paper, we propose a graph model for representing complex regions, which overcomes the above shortcomings.

For a complex region A, we decompose both the *interior* and the *exterior* of A into connected components. To avoid exotic situations, we suppose the closure of each bounded component is a simple region with holes. We then represent each complex region as a link graph, where a node represents a connected component of the interior or the exterior of the complex region, and two components are linked if their boundary share a curve. We also define a natural level function for components of a complex region such that any two linked components have successive levels.

For a srh A, let $\widehat{A}$ be the union of the srh and its holes. Then $\widehat{A}$, called the *generalized region* of A, is also a simple region (Egenhofer, Clementini, and Di Felice 1994). Given the link graph, the holes and the generalized region of each bounded component can be computed efficiently. As for the unbounded component, we prove that its complement is actually a composite region that consists of finitely many simple regions. Note that the generalized region and holes of a bounded component are simple regions. We call these simple regions the *atoms* of the complex region.

We prove that these atoms are necessary and complete for locally determining the nine-intersections between the interiors, the boundaries, and the exteriors of two complex regions (the 9IM relation henceforth). The 9IM relations are identically the same as the RCC8 basic relations if only simple regions are concerned. This means, on one hand, knowing the RCC8 relations between the atoms of two complex regions is sufficient to determine the 9IM relation between these complex regions; on the other hand, if one atom is removed from the atom set of a complex region A, there exist two complex regions B_1, B_2 such that the 9IM relation between A and B_1 is different from that between A and B_2, despite that all corresponding local RCC8 relations are equal.



(a) (b)

Figure 3: Two complex regions with the same W-B tree

Our graph representation provides a natural method for comparing two complex regions. It also gives rise to a natural method for generalizing complex regions step by step. This introduces a new idea for map generalization, which is a very important technique used in cartography and GISs.

The rest of this paper proceeds as follows. Section 2 recalls some basic notions and preliminary results about complex regions, and Section 3 proposes the graph model. A complete characterization of the (*global*) 9IM relations between two complex regions *locally* is given in Section 4, and the generalization technique is introduced in Section 5. The last section concludes the paper.

Backgrounds

In this section, we introduce the basic notions and preliminary results needed in this paper.

Basic topological notions

A *topology* $\mathcal{T}$ over a nonempty set U is a subset of the powerset $\wp(X)$ that is closed under arbitrary unions and finite intersections. We call $(X, \mathcal{T})$ a *topological space*, and call each set in $\mathcal{T}$ an *open set*. For a set A of X, the *interior* of A, denoted by A°, is the largest open set which is contained in A.

Given a topological space $(X, \mathcal{T})$, a set $A \subseteq X$ is a *closed set* if its complement $X \setminus A$ is open. For any $B \subseteq X$, the *closure* of B is the smallest closed set, written $\overline{B}$, which contains B. A closed set A is *regular* if $\overline{A^\circ} = A$.

The *boundary* of a subset A of X, denoted by ∂A, is defined to be the set difference of $\overline{A}$ and A°, *i.e.* $\partial A = \overline{A} \setminus A^\circ$.

A subset A of a topological space $(X, \mathcal{T})$ is *disconnected* if there exist two disjoint open sets U, V such that $U \cap A$ and $V \cap A$ are nonempty and $A \subseteq U \cup V$. We say A is a *connected set* if it is not disconnected.

A connected set U is called a *(connected) component* of an open set A if U is a maximal connected subset of A. Two different connected components are clearly disjoint.

Complex regions and components

In this paper, we concern ourselves with areal objects in the real plane (with the usual topology). As usual, we define a *plane region* to be a regular closed set in the real plane.

For a bounded plane region A, we call the closure of a component of A° (the interior of A) a *positive* component of A, and call the closure of a component of A^e (the exterior of A) a *negative* component of A. Since A is bounded, A^e has a unique unbounded component. We call its closure *the* unbounded component of A and denote the closure as b_0. We call the closure of each bounded component of A^e a *hole* component of A, or simply a *hole* of A. As a consequence, each component of A is a plane region, *i.e.* a regular closed set in the plane.

Clearly, each bounded region has a unique unbounded component and at least one positive component. It is also easy to see that a bounded region may have an infinite number of positive and/or hole components. For practical applications, it is reasonable to require a complex region to have only finitely many (positive or hole) components.

In this paper, we plan to establish a valid representation scheme for plane regions. A primary requirement for such a representation is that *each region is finitely representable*. We hope each object can be reconstructed by applying finite basic operations (*e.g.* union, intersection, and difference) upon a finite set of atomic regions. For our purpose, it is reasonable to choose simple regions as atomic regions.

A complex region is defined as the following.

Definition 1 (complex region)**.** *A complex region A is a bounded regular closed subset of the real plane that has a finite set of positive components and a finite set of holes, where each positive component of A and each hole of A is a simple region with holes.*

For a bounded component c, suppose $a_1, a_2, \cdots, a_k$ are its holes. Let $a_0 = c \cup \bigcup_{i=1}^k a_i$. We call a_0 the *generalized region* of c. To avoid exotic situations, we assume the generalized region and the holes of a bounded component (which is a srh) are all simple regions. For clarity, we often write c as $(a_0; a_1, \cdots, a_k)$ and $\widehat{c}$ for the generalized region a_0 of c.

In the following section, we will propose a graph representation for complex regions, and then show how to compute the generalized region and the holes of a component.

A graph representation of complex region

Our layered graph representation of complex regions is based on components and strong connectedness.

Link graph

We recall that a simple curve in the plane is a subset of the real plane that is topologically equivalent to the closed interval $[0, 1]$. The boundary of each simple region contains a simple curve.

Proposition 1. *For a complex region, the intersection of two positive (negative, resp.) components is a finite set. The intersection of a positive and negative component is a finite set or contains a simple curve.*

We say two components are *strongly connected* or *linked* if the intersection of their closures contains a simple curve. Each component is linked with at least one other component. This is because its boundary is contained in the union of the boundaries of all other components.

For a complex region A, we introduce a level function for the components of A. For each component c of A, we define $\mathsf{lev}(c)$, the level of component c, inductively as follows:



Figure 4: The link graphs of regions in Fig. 1



Figure 5: The link graphs of regions in Fig. 3

- The level of b_0, the unbounded component of A, is 0;
- For an undefined c^*, if there exists a previously defined node c which is linked to c^*, then define $\mathsf{lev}(c^*) = \mathsf{lev}(c) + 1$.

For two linked components c_1, c_2, it is easy to prove that $\mathsf{lev}(c_1) - \mathsf{lev}(c_2) = \pm 1$. Take the complex region in Figure 3(b) as example. It is clear that $\mathsf{lev}(b_0) = 0$, $\mathsf{lev}(a_1) = \mathsf{lev}(a_2) = 1$, and $\mathsf{lev}(b_1) = 2$.

Definition 2 (link graph)**.** *The link graph $\mathcal{L}_A$ of a complex region A is defined as a directed graph $(N(A), E(A))$ as follows:*

- *$N(A)$ is the set of all components (positive, hole, or unbounded) of A;*
- *For $c_1, c_2 \in N(A)$, $(c_1, c_2) \in E(A)$ if they are linked and $\mathsf{lev}(c_2) = \mathsf{lev}(c_1) + 1$.*

Note that if two nodes are linked, then exactly one is positive. If there is a directed edge from c_1 to c_2, then we call c_1 a *parent* of c_2, and call c_2 a *child* of c_1. We say a component c is a *leaf* if it does not have any children.

In this way, we represent each complex region by a layered graph. Fig.s 4 and 5 show the link graphs for the regions in Fig.s 1 and 3, respectively. Each pair of regions in these two figures have different link graphs. This suggests that link graph model is more expressive than the models proposed in (Schneider and Behr 2006) and (Worboys and Bofakos 1993).

We next use link graph to characterize simple regions with holes and composite regions (Clementini, Di Felice, and Califano 1995).

Proposition 2. *Let A be a complex region. Then A is a simple region with holes iff the link graph $\mathcal{L}_A$ has a unique node at level 1 and has no node with level greater than 2.*



Figure 6: The link graph of a srh with k-holes



Figure 7: The link graph of a composite region with k-atoms

In other words, a complex region is a srh iff its link graph has the form as the one given in Fig. 6

A complex region is called a *composite region* if its positive components are all simple regions and it has no hole components. The following characterization is clear.

Proposition 3. *A complex region is a composite region iff its link graph has no node with level above 1.*

In other words, a complex region is a composite region iff its link graph has the form as the one given in Fig. 7

Computing the holes and the generalized region

Suppose the link graph $\mathcal{L}_A$ of a complex region A is given. The following questions are natural. For a node n_1 in $\mathcal{L}_A$, *is the component represented by* n_1 *a simple region*? and if not, *how to compute its generalized region and holes*?

The following proposition shows that a component has holes iff the corresponding node separates the link graph.

Proposition 4. *Suppose c is a bounded component of a complex region A. Then c has no holes iff the (undirected) graph obtained by removing c from* $\mathcal{L}_A$ *is still connected.*

Proof. If h is a hole of c, then h is the union of several components of A. Suppose h contains a component c_1. Then it is easy to see that any path from the root b_0 to c_1 must pass through c. This means $\mathcal{L}_A$ will be disconnected if removing c from it. On the other hand, if h has no hole, then for any descendant c_1 of c there is another path from b_0 to c_1. That is, $\mathcal{L}_A$ is still connected if removing c from it. $\square$

Example 1. *Figure 8 shows a much complicated region and its link graph, where an arrow from* c_1 *to* c_2 *suggests that the level of* c_1 *is lower than that of* c_2*. For this complex region, it is easy to see that only* a_2 *has a hole, the other bounded components are simple regions.*



Figure 8: A complex region A and its link graph

For a node (*i.e.* a component) c, we next compute its generalized region and holes. Suppose the link graph $\mathcal{L}_A$ is decomposed into $k+1$ connected subgraphs $\mathcal{L}_0(c), \mathcal{L}_1(c), \cdots, \mathcal{L}_k(c)$ by removing node c from $\mathcal{L}_A$, where $\mathcal{L}_0(c)$ is the connected subgraph that contains b_0.

Theorem 1. *Suppose* $c \neq b_0$*. For each* $i \geq 1$*, the union of the components in* $\mathcal{L}_i(c)$ *is a hole of c, and the generalized region of c is the union of c and all its holes.*

Proof. Note that if c_1 is not a descendant of c, then there is a path in $\mathcal{L}_A$ from b_0 to c_1 that does not pass through c. This means, c_1 is a node in $\mathcal{L}_0(c)$. So, all non-descendants of c are nodes in $\mathcal{L}_0(c)$. For each $i \geq 1$, the union of all nodes (*i.e.* components) in $\mathcal{L}_i(c)$ is a simple region h_i. Write $\widehat{c}$ for the generalized region of c, *i.e.* the union of c and all h_i. Then $\widehat{c}$ is also a simple region, and c is the simple region with holes $(\widehat{c}; h_1, \cdots, h_k)$. $\square$

The above theorem describes how to compute the holes and the generalized region of a component. The components that are contained in these holes can be characterized as follows.

Proposition 5. *For c,* $\mathcal{L}_i(c)$ *as in the above theorem, a descendant* c_1 *of c is in* $\mathcal{L}_i(c)$ *for some* $i \geq 1$ *iff each path from* b_0 *to* c_1 *passes through c.*

If the unbounded component b_0 itself is removed from $\mathcal{L}_A$, then $\mathcal{L}_0(c) = \varnothing$. In this case, the union of all components in a connected subgraph $\mathcal{L}_i(c)$ is also a simple region, written b_i. It is straightforward to see that $-b_0$, the closure of the complement of b_0, is the union of all these b_i. Clearly, $-b_0$ has no holes and b_0 is its unbounded component. This implies that $-b_0$ is a composite region.

Proposition 6. *Let b_0 be the unbounded component of a complex region A. Then $-b_0$, the closure of the complement of b_0 is a composite region.*

Recall that the generalized region of a srh is the union of all its bounded components. For a complex region A, it is reasonable to call the union of all its bounded components the generalized region of A. Write $\widehat{A}$ for the generalized region of A. By Prop. 6, we know $\widehat{A}$ is the composite region $-b_0$.

We now define the atom set of a complex region.

Definition 3 (atom set of complex region)**.** *A simple region a is called an atom of a complex region A if one of the following conditions is satisfied:*

- *there exists a bounded component c of A such that a is the generalized region or a hole of c;*
- *a is a positive component of the composite region $-b_0$, where b_0 is the unbounded component of A.*

We write $\mathit{ATOM}(A)$ for the atom set of A.

Local characterization of the 9IM relations

For a bounded complex region A, we write b_0 for its unbounded component, and write $\Gamma^+(A)$ and $\Gamma^-(A)$ for the sets of bounded positive and hole components of A, respectively. Let $\Gamma(A) = \Gamma^+(A) \cup \Gamma^-(A)$ for the set of bounded components of A. Note $b_0 \notin \Gamma(A)$.

For two complex regions A and A', the 9IM relation (Egenhofer and Franzosa 1991) between A and A' is defined as

$$M(A, A') = \left(\begin{array}{ccc} A^\circ \cap A'^\circ & A^\circ \cap \partial A' & A^\circ \cap A'^e \\ \partial A \cap A'^\circ & \partial A \cap \partial A' & \partial A \cap A'^e \\ A^e \cap A'^\circ & A^e \cap \partial A' & A^e \cap A'^e \end{array} \right).$$

There are eight different 9IM relations between simple regions. These are identically the RCC8 basic relation. It has also been proved that there are all together 34 different 9IM relations between (bounded) complex regions (Schneider and Behr 2006; Li 2006).

In this section, we characterize the 9IM relations in terms of the components of the two complex regions.

We introduce the following notion of intersection degree.

Definition 4 (intersection degree)**.** *The positive intersection degree of a component c of A, w.r.t. A', denoted by $\deg^+_{A'}(c)$ or simply $\deg^+(c)$, is the number of positive components a' of A' such that $c^\circ \cap A'^\circ \neq \varnothing$,* i.e.

$$\deg^+(c) = |\{a' \in \Gamma^+(A') : c^\circ \cap A'^\circ \neq \varnothing\}|. \qquad (1)$$

Similarly, the intersection degree of c, w.r.t. A', denoted by $\deg_{A'}(c)$ or simply $\deg(c)$, is defined as

$$\deg(c) = |\{d' \in \Gamma(A') \cup \{b'_0\} : c^\circ \cap d'^\circ \neq \varnothing\}|, \qquad (2)$$

where b'_0 is the unbounded component of A'.

We next give each of the nine intersections a characterization in terms of components. For convenience, we write, say, $I_{\circ\partial}$, for the intersection of A° and $\partial A'$. Note that I_{ee} is always nonempty because A and A' are bounded. For $I_{\partial\partial}$ we have

Lemma 1 ($I_{\partial\partial}$)**.** $\partial A \cap \partial A' = \varnothing$ iff $\partial a \cap \partial a' = \varnothing$ *for all* $a \in \Gamma^+(A)$ *and all* $a' \in \Gamma^+(A')$.

Proof. This is because $\partial A = \bigcup\{\partial a : a \in \Gamma^+(A)\}$ and $\partial A' = \bigcup\{\partial a' : a' \in \Gamma^+(A')\}$. ☐

The remaining seven intersections can be determined by the intersection degrees given in Dfn. 4.

Lemma 2 ($I_{\circ\circ}$)**.** $A^\circ \cap A'^\circ \neq \varnothing$ iff $\deg^+(a) > 0$ *for some* $a \in \Gamma^+(A)$.

Lemma 3 ($I_{\circ\partial}$)**.** $A^\circ \cap \partial A' \neq \varnothing$ iff $\deg(a) > 1$ *for some* $a \in \Gamma^+(A)$.

The case of $I_{\partial\circ}$ is similar.

Lemma 4 ($I_{\circ e}$)**.** $A^\circ \cap A'^e \neq \varnothing$ iff $\deg^+(b'_0) > 0$ *or* $\deg^+(b') > 0$ *for some* $b' \in \Gamma^-(A')$.

The case of $I_{e\circ}$ is similar.

Lemma 5 ($I_{\partial e}$)**.** $\partial A \cap A'^e \neq \varnothing$ iff $\deg(b'_0) > 1$ *or* $\deg(b') > 1$ *for some* $b' \in \Gamma^-(A')$.

The case of $I_{e\partial}$ is similar.

We take the proof of the case $I_{\circ\partial}$ as example.

Proof. $A^\circ \cap \partial A' \neq \varnothing$ iff there exist $a \in \Gamma^+(A)$ such that $a^\circ \cap \partial A' \neq \varnothing$. The latter is equivalent to saying that there exist a positive A'-component a' and a negative A'-component b' such that $a^\circ \cap A'^\circ \neq \varnothing$ and $a^\circ \cap b'^\circ \neq \varnothing$. This is also equivalent to $\deg(a) > 1$. ☐

In the following, we consider how to determine the nine intersections locally, *i.e.* by using atoms of A and A'. We begin with $I_{\partial\partial}$. By Lemma 1, we need check if $\partial a \cap \partial a' = \varnothing$ for all positive components $a \in \Gamma^+(A), a' \in \Gamma^+(A')$. Because a bounded component is a srh, we need only consider the intersection of the boundaries of two srhs.

Lemma 6. *Let $A = (a_0; a_1, \cdots, a_k)$ and $A' = (a'_0; a'_1, \cdots, a'_l)$ be two srhs. Then $\partial A \cap \partial A' \neq \varnothing$ iff $\partial a_i \cap \partial a'_j \neq \varnothing$ for some $0 \leq i \leq k$ and some $0 \leq j \leq l$.*

Proof. This is because the boundary of a srh is the union of the boundaries of all its simple regions. ☐

Therefore, the boundary-boundary intersection of two complex regions can be reduced to the boundary-boundary intersection of atoms. To determine the other intersections, we check the following two types of intersections:

I1. Is $c^\circ \cap c'^\circ$ empty for two bounded components c, c'?

I2. Is $c^\circ \cap {b'_0}^\circ$ empty for a bounded component c and an unbounded component b'_0?

The intersection of two bounded components can be decided in the following way.

Lemma 7. *Let $A = (a_0; a_1, \cdots, a_k)$ and $A' = (a'_0; a'_1, \cdots, a'_l)$ be two srhs. Then $A^\circ \cap A'^\circ = \varnothing$ iff $a_0^\circ \cap {a'_0}^\circ = \varnothing$, or $a_0 \subseteq a'_i$ for some i, or $a'_0 \subseteq a_j$ for some j.*

For the unbounded component b_0 of a complex region, by Prop. 6 we know $-b_0$, the closure of the set complement of b_0, is a composite region, *i.e.* $-b_0$ has no negative components and all its positive components are simple regions.

Lemma 8. *Let $A = (a_0; a_1, \cdots, a_k)$ be a simple region with holes. Assume that $-b_0$ is composed of l simple regions $b_1, b_2, \cdots, b_l$. Then $A^\circ \cap b_0^\circ \neq \varnothing$ iff a_0 is contained in some b_i $(1 \leq i \leq l)$.*

Proof. This is because $A^\circ \cap b_0^\circ = \varnothing$ iff $A \subset -b_0$. Since A° is a connected set, it must be contained in a component of $-b_0$. That is, A°, hence A, must be contained in some simple region b_i, which is equivalent to saying that the generalized (simple) region of A, *viz.* a_0, is contained in b_i. ☐

By the above lemmas, we know the nine intersections of two complex region A, A' can be determined by the RCC8 (or equivalently 9IM) relations between atoms of A and A'.

Theorem 2 (sufficiency)**.** *The 9IM relation between two complex regions A, A' can be uniquely decided by the RCC8 relations between the atoms of A and A'.*

Are these atoms also *necessary* for determining the 9IM relation between A and A'? That is, if we delete an atom o from the atom set $\mathsf{ATOM}(A)$, are there two complex regions $B \neq C$, such that $M(A,B) \neq M(A,C)$ but $M(\rho,B) = M(\rho,C)$ for all $\rho \in \mathsf{ATOM}(A)$ such that $\rho \neq o$?

The answer is positive. To show an atom o of A is necessary, we show there is a simple region o' such that $M(A,o) \neq M(A,o')$ but $M(\rho,o) = M(\rho,o')$ for all A-atoms $\rho \neq o$. This means, as far as 9IM relations are concerned, o' behaves identically as o locally, but different globally.

We illustrate the construction of o' by one example. Consider the complex region A given in Figure 9(a). Let b be the union of a_1, a_2 and b_1. Then b is a simple region, and the atom set of A is $\{a_1, a_2, b_1, b\}$. We show that b_1 and b are necessary. Let b' be the simple region as shown in Figure 9(b). Then b' has the same RCC8 relations with a_1, a_2 and b as b_1 does. But $M(A,b_1) \neq M(A,b')$, because $\partial b_1 \subset \partial A$ while $\partial b' \not\subseteq \partial A$. As for b, take a simple region b^* that contains b (see Figure 9(c)). Then b and b^* have the same 9IM relations with a_1, a_2 and b_1. But since $\partial b \subset \partial A$ and $\partial b^* \not\subseteq \partial A$, we know $M(b,A) \neq M(b^*,A)$.

In general, we have

Lemma 9. *Suppose A is a complex region and o is an atom of A. Then there exists another simple region o' such that $M(A,o) \neq M(A,o')$ but $M(\rho,o) = M(\rho,o')$ for all atoms $\rho \neq o$ of A.*

Proof. Note that $o \in \mathsf{ATOM}(A)$ implies $\partial o \subseteq \partial A$. We construct a simple region o' that is similar to o but $\partial o' \not\subseteq \partial A$.

We take the case when o is the generalized region of a bounded component c as example. For each $\rho \in \mathsf{ATOM}(A)$, take a point P_ρ in $\partial o \cap \partial \rho$ if it is nonempty. Note that there exists at least a segment α of the closed Jordan curve ∂o that is disjoint from all the holes of c. Take a point $P \in \alpha$ such that $P \neq P_\rho$ for all ρ. So the distance from P to the holes of c is nonzero. Removing a tiny part around P from o such that the resulting region is still simply connected and disjoint from all holes of o. Write o' for this simple region. Clearly, there is a point Q in the boundary of o' which is not contained in the boundary of A. This implies $M(A,o) \neq M(A,o')$. On the other hand, for each A-atom $\rho \neq o$, if $\partial\rho \cap \partial o$ is empty, so is $\partial\rho \cap \partial o'$; and if $\partial\rho \cap \partial o$ is nonempty, then $P_\rho \in \partial\rho \cap \partial o' \neq \varnothing$. Moreover, the part-whole relation between o' and each ρ is the same as that between o and ρ. Therefore, $M(\rho,o) = M(\rho,o')$ for all atoms $\rho \neq o$ of A.



(a) (b) (c)

Figure 9: Three complex regions

The case when o is a hole of a bounded component c or when o is a component of $-b_0$ is similar. ☐

As a consequence, we know each atom in $\mathsf{ATOM}(A)$ is necessary.

Theorem 3 (necessary)**.** *Each atom of a complex region A is necessary in locally determining the 9IM relation of A and some other region.*

A special case has been investigated by (McKenney, Praing, and Schneider 2008). Note that if $A = (a_0; a_1, \cdots, a_k)$ is a srh, then $-b_0 = a_0$.

Corollary 1. *Let $A = (a_0; a_1, \cdots, a_k)$ and $A' = (a_0; a'_1, \cdots, a'_l)$ be two srhs. Then the 9IM relation between A and A' is completely determined by the RCC8 relations between all a_i and all a'_j.*

Generalization by dropping

Map generalization is a very important technique used in cartography and GISs. We now propose a new method to generalize a complex region into simpler ones.

For a simple region with holes $A = (a_0; a_1, \cdots, a_k)$, we obtain its generalized region $\widehat{A}$ by dropping all its holes. Similarly, for a complex region, we obtain its generalized region $\widehat{A}$ by merging all its bounded components. We could also obtain less complicated regions by dropping *some* components together with their holes from a complex region. Take Figure 8 as example, b_2 is the unique hole of a_2. Dropping b_2 could obtain a simpler complex region.

We next generalize a complex region step by step by dropping one component at a time.

Let A be a complex region with link graph $\mathcal{L}_A$. Suppose c is a component that has no children. Let $p_1, \cdots, p_k$ be the parents of c. It is easy to see that $r = \bigcup\{p_i\}_{i=1}^k \cup \{c\}$



Figure 10: A complex region (upper) and its generalization (lower), where $b_1^* = a_2 \cup b_1 \cup b_2 \cup b_3$.

is also a connected region. In this way, we obtain a new complex region A_1 by merging the hole c into its parents. Note that each A-component other than c and $p_1, \cdots, p_k$ is an A_1-component. Note r is also a component of A_1. The link graph of A_1 is obtained by setting

- $V_1 = (V \setminus \{c, p_1, \cdots, p_k\}) \cup \{r\}$;
- $E_1 = (E \cap V_1 \times V_1) \cup \{(m, r) : (\exists i)(m, p_i) \in E\}$.

Figure 10 shows a complex region and one step of its generalization.

Note that after each step of dropping, we obtain a complex region that has fewer components. Step by step, we will obtain that a region that cannot be generalized. This is exactly $\widehat{A}$, the *generalized region* of A.

Conclusion

In this paper we have established a qualitative model for representing the internal structure of complex plane regions. For the first time, we introduced the notions of holes, generalized regions, and atoms to general complex regions. In particular, we define the generalized region of a complex region as the the union of all bounded components. We have proved that the generalized region of a complex region is a composite region and equals to the closure of the set complement of the unbounded component.

In the layered graph of a complex region A, a node represents a component of the complex region, and an edge exists between two nodes iff the two components share a common one-dimensional boundary. Based on this model, we gave methods for computing the atoms of A, which are either holes or the generalized region of a bounded component of A, or the atoms of the generalized region of A.

We have proved that these atoms are necessary and sufficient for determining the nine-intersection relation between complex regions. We believe this provides a partial justification for the rationality for applying the 9IM approach on complex region. It also suggests that the 9IM relation between complex regions can be implemented through the implementation of the RCC8 (or the 9IM) relation between simple regions.

Furthermore, the hierarchical representation also provides a natural way for generalizing the complex region. This is very important in automatic cartography and GISs.

Future work will consider spatial reasoning with this internal structure model. Some related work has been done for simple regions with holes (Vasardani and Egenhofer 2009).

References

[Clementini, Di Felice, and Califano 1995] Clementini, E.; Di Felice, P.; and Califano, G. 1995. Composite regions in topological queries. *Information Systems* 20(7):579–594.

[Cohn and Hazarika 2001] Cohn, A., and Hazarika, S. 2001. Qualitative spatial representation and reasoning: An overview. *Fundamenta Informaticae* 46(1-2):1–29.

[Cohn and Renz 2007] Cohn, A., and Renz, J. 2007. Qualitative spatial reasoning. In van Harmelen, F.; Lifschitz, V.; and Porter, B., eds., *Handbook of Knowledge Representation*. Elsevier.

[Egenhofer and Franzosa 1991] Egenhofer, M., and Franzosa, R. 1991. Point-set topological spatial relations. *International Journal of Geographical Information Systems* 5(1):161–174.

[Egenhofer, Clementini, and Di Felice 1994] Egenhofer, M.; Clementini, E.; and Di Felice, P. 1994. Topological relations between regions with holes. *International Journal of Geographical Information Systems* 8(2):129–142.

[Kontchakov et al. 2008] Kontchakov, R.; Pratt-Hartmann, I.; Wolter, F.; and Zakharyaschev, M. 2008. Topology, connectedness, and modal logic. In Areces, C., and Goldblatt, R., eds., *Advances in Modal Logic*, 151–176. College Publications.

[Li 2006] Li, S. 2006. A complete classification of topological relations using the 9-intersection method. *International Journal of Geographical Information Science* 20(6):589–610.

[McKenney, Praing, and Schneider 2008] McKenney, M.; Praing, R.; and Schneider, M. 2008. Deriving topological relationships between simple regions with holes. In *SDH*, 1863–2246.

[Randell, Cui, and Cohn 1992] Randell, D.; Cui, Z.; and Cohn, A. 1992. A spatial logic based on regions and connection. In *Proceedings of the Third International Conference on Principles of Knowledge Representation and Reasoning (KR-92)*, 165–176.

[Schneider and Behr 2006] Schneider, M., and Behr, T. 2006. Topological relationships between complex spatial objects. *ACM Trans. Database Syst.* 31(1):39–81.

[Varzi 2007] Varzi, A. 2007. Spatial reasoning and ontology: parts, wholes, and locations. In Aiello, M., P.-H. I. v. B. J., ed., *Handbook of Spatial Logics*. Springer.

[Vasardani and Egenhofer 2009] Vasardani, M., and Egenhofer, M. 2009. Comparing relations with a multi-holed region. In *Proceedings of the 9th International Conference on Spatial Information Theory (COSIT 2009)*, 159–176.

[Worboys and Bofakos 1993] Worboys, M., and Bofakos, P. 1993. A canonical model for a class of areal spatial objects. In *Proceedings of the Third International Symposium on Advances in Spatial Databases (SSD'93)*, 36–52. London, UK: Springer-Verlag.